

READ SAFETY DIRECTIONS BEFORE OPENING OR USING

# **Moncut**<sup>®</sup> SC Fungicide

Active Constituent: 464 g/L FLUTOLANIL


GROUP **G** FUNGICIDE

**For the control of seed-borne  
black scurf (*Rhizoctonia solani*)  
in potatoes as per the  
DIRECTIONS FOR USE Table.**

**IMPORTANT:** READ THE ATTACHED BOOKLET BEFORE USE

**Net Contents: 5L**

Certis Australia Pty. Limited A.B.N. 28 100 543 986  
Level 11, Tower B, Zenith Centre,  
821 Pacific Highway, Chatswood NSW 2067  
Phone: (02) 8448 8155 Fax: (02) 8448 8156



READ SAFETY DIRECTIONS BEFORE OPENING OR USING

# MONCUT SC FUNGICIDE

Active Constituent: 464 g/L FLUTOLANIL

For the control of seed-borne black scurf (*Rhizoctonia solani*) in potatoes as per the DIRECTIONS FOR USE Table.

## STORAGE AND DISPOSAL

### Product and product container

Keep out of reach of children.

Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight.

Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank when diluting product. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

### Application equipment

Thoroughly rinse all application equipment with clean water immediately after use and dispose of rinsings in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots.

## SAFETY DIRECTIONS

Will irritate the eyes. Avoid contact with eyes. Wash hands after use.

### FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre (telephone 13 11 26).

### MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet available from Certis Australia Pty. Limited.

### EXCLUSION OF LIABILITY

This product as supplied is of a high grade and suitable for the purpose for which it is expressly intended and must be used in accordance with the directions. The user must monitor the performance of any product as climatic, geographical or biological variables and/or developed resistance may affect the results obtained. No responsibility is accepted in respect of this product, save for those non-excludable conditions implied by the Trade Practices Act or any State or Federal legislation.

NRA Approval No.: 53217 / 0403

Moncut® is a Registered Trademark of Nihon Nohyaku Co. Ltd.

**IMPORTANT: READ THIS BOOKLET BEFORE USE**

## DIRECTIONS FOR USE

CROP	DISEASE	RATE	CRITICAL COMMENTS
Potatoes	Black scurf ( <i>Rhizoctonia solani</i> ) (seed-borne infections only)	150 mL/ tonne of seed potatoes	Spray Moncut onto seed potatoes at or prior to planting. Prior to application Moncut should be diluted with sufficient water to ensure thorough, even coverage of seed potatoes. A minimum total application volume of 3 litres per tonne of seed is recommended. Where conditions are likely to favour seed piece decay (e.g. high temperatures and wet soil soon after planting): <ul style="list-style-type: none"><li>• Ensure cut tubers are properly cured before application</li><li>• Ensure treated seed is dry prior to planting</li><li>• Avoid treatment on the planter, or immediately prior to planting.</li></ul>

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

**WITHOLDING PERIOD NOT REQUIRED WHEN USED AS DIRECTED.**

## GENERAL INSTRUCTIONS

Moncut SC Fungicide is for the protection of potatoes against seed-borne black scurf (*Rhizoctonia solani*). Moncut is sprayed onto seed potatoes prior to planting, or at the time of planting. When applied as directed, this product will protect progeny tubers from infection via seed-borne *Rhizoctonia solani*. No claim is made for protection of tubers against soil-borne *Rhizoctonia solani*. Best results will be achieved where other management practices (such as crop rotation) are employed to reduce the occurrence of soil-borne *Rhizoctonia solani*.

## GROUP **G** FUNGICIDE

### Fungicide Resistance Warning

Moncut is an anilide fungicide. For fungicide resistance management, Moncut is a Group **G** fungicide. Some naturally-occurring fungal biotypes resistant to Moncut, and other Group **G** fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungal population. The resistant individuals can eventually dominate the fungal population if these fungicides are used repeatedly. These resistant fungi will not be controlled by Moncut and other Group **G** fungicides, thus resulting in a reduction in efficacy and possible yield loss.

Since the occurrence of resistant fungi is difficult to detect prior to use, Certis Australia Pty Ltd accepts no liability for any losses that may result from the failure of Moncut to control resistant fungi.

### Export of Treated Produce

Growers should note that suitable MRLs or import tolerances may not be established in all markets for produce treated with Moncut SC Fungicide. If you are growing produce for export, please check with Certis Australia Pty Ltd for the latest information on MRLs and import tolerances **BEFORE** using Moncut.

### Equipment

Moncut is applied to seed potatoes prior to, or at planting. Application must be conducted with an applicator suitable for liquid treatments. The equipment must be accurately calibrated to ensure even coverage and correct application rates.

### Application

Prior to application Moncut should be diluted with sufficient water to ensure thorough, even coverage of seed potatoes. A minimum total application volume of 3 litres per tonne seed potatoes is recommended. Apply the diluted product onto dry tubers with a fine spray, ensuring even coverage of all surfaces. Tubers should be free from soil deposits at application.

Diluted product should be constantly agitated in the spray tank during application. Prepare sufficient product for the quantity of seed to be treated. Do not store diluted product.

### Compatibility

If using dust treatments (firbark, lime or mancozeb), these should be applied to tubers before applying Moncut. The use of cement dust is not recommended on tubers to be treated with Moncut.

## PRECAUTIONS

Do not use treated seed potatoes for human or animal consumption.

Do not allow potatoes intended for human or animal consumption to become contaminated with this product.

Store treated seed potatoes away from other potatoes, animal feed or foodstuffs, in a cool, dry storage area. Clearly label bins, bags or other containers of treated seed potatoes, to indicate that they have been treated with Moncut. Do not use bins, bags or containers that have held Moncut treated potatoes for any other purpose, including storage of potatoes intended for human or animal consumption.

## PROTECTION OF LIVESTOCK

**DO NOT** use treated potato seed for animal consumption.

**DO NOT** allow feed intended for animal consumption to become contaminated with this product.

## PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

**DO NOT** contaminate streams, rivers or waterways with the product or used containers.