

# CAUTION

**KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

# PAYBACK<sup>®</sup>

## Plant Growth Regulator

**ACTIVE CONSTITUENT: 250 g/L PACLOBUTRAZOL**

To reduce vegetative growth in mango, stone fruit and apple trees.

**IMPORTANT: READ THE ATTACHED LEAFLET BEFORE USING THIS PRODUCT**

## CONTENTS: 1 & 5 LITRES

Crop Care Australasia Pty Ltd, 77 Tingira Street, Pinkenba, Queensland 4008

**STORAGE AND DISPOSAL**

Store in the closed, original container in a cool, well ventilated area. Do not store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

**SAFETY DIRECTIONS**

Avoid contact with eyes and skin. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water.

**FIRST AID**

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

**MATERIAL SAFETY DATA SHEET**

Additional information is listed in the Material Safety Data Sheet.

**Conditions of sale**

Crop Care Australasia Pty Ltd will not accept any responsibility whatsoever and howsoever arising and whether for consequential loss or otherwise in connection with the supply or use of these goods other than responsibility for the merchantable quality of the goods and such responsibilities mandatorily imposed by Statutes applicable to the sale or supply of these goods. To the extent allowed by such Statutes the liability of Crop Care Australasia Pty Ltd is limited to the replacement of the goods or (at the option of Crop Care Australasia Pty Ltd) the refund of the price paid and is conditional upon a claim being made in writing and where possible sufficient part of the goods to enable proper examination being returned to Crop Care Australasia Pty Ltd within thirty days of delivery.

In a Transport Emergency Dial 000 Police or Fire Brigade	<b>SPECIALIST ADVICE IN EMERGENCY ONLY</b> 1800 033 498 <b>ALL HOURS – AUSTRALIA WIDE</b>
--	---

Crop Care Australasia Pty Ltd is the licensed user of Payback which is a Registered Trademark.



Crop Care Australasia Pty Ltd  
ACN 061 362 347  
77 Tingira Street  
PINKENBA Qld 4008

Insert barcode  
and label code

APVMA Approval No. : 58627/1L/0404; 58627/5L/0404

**BN :**

**DOM :**

# CAUTION

**KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

# PAYBACK<sup>®</sup>

## Plant Growth Regulator

**ACTIVE CONSTITUENT: 250 g/L PACLOBUTRAZOL**

To reduce vegetative growth in mango, stone fruit and apple trees.

**IMPORTANT: READ THIS LEAFLET BEFORE USING THIS PRODUCT**

Crop Care Australasia Pty Ltd, 77 Tingira Street, Pinkenba, Queensland 4008

## DIRECTIONS FOR USE

DO NOT use on trees of low vigour or under stress (see General Instructions).

DO NOT use in orchards underplanted with any other crops.

DO NOT apply to young trees which have recently been interplanted in orchards and which are required to make vigorous growth.

DO NOT apply by aircraft.

Crop & Situation	State	Application Method	Rate	Critical Comments
<b>Mango</b>	Qld, NT & NSW only	Collar, Drench: <u>Trees 3 - 6 yrs old:</u> Light sands Clay, red volcanics  <u>Trees over 6 yrs:</u> Light sands Clay, red volcanics	Per tree: 8 to 12mL 12 to 16mL  Per tree: 16 to 20mL 20mL	Apply within 4 weeks after harvest or no later than mid February. Adjust rate within range according to tree age. Use high rate for older trees. Ensure trees are well watered before and after application. <b>CAUTION:</b> Do not treat young trees with a canopy of less than 3m diameter. <b>CAUTION:</b> Excessive compaction of flower panicles may result from application at times later than recommended, or from delayed uptake caused by dry soil conditions. Care should be taken to ensure correct insecticide application to control webworm in compact flower panicles. <b>RETREATMENT:</b> The effects of 'Payback' on flowering and growth may persist for more than one season. <b>Sandy soils:</b> Postpone retreatment until after the harvest following resumption of normal growth. Use the interval between the first treatment and resumption of normal growth as a guide to retreatment frequency in subsequent years. <b>Clays &amp; Red Volcanic Soils:</b> Retreat each season. If reduction of growth increases progressively from season to season postpone treatment until after the harvest following the resumption of normal growth.
<b>Apples - Red Delicious &amp; Granny Smith</b>	All States	Spray	500mL or 1L/ha Add 18mL Shirwet 600 or Nufarm Surfactant 600 per 100L water in final spray volume	Apply in a sequential spray program commencing 28 days after full bloom on Red Delicious or 35 days after full bloom on Granny Smith. Use lower rate on less vigorous trees. Apply subsequent sprays at the lower rate at intervals of 21 days for as long as growth control is required. Activity results from uptake of chemical by young green stem tissue. Do not apply more than 4L/ha per season. <b>CAUTION:</b> If using Carbaryl as a chemical thinning agent, plan to allow 21 days between application of Carbaryl and Payback.
<b>Peaches Nectarines</b>	All States	Trickle irrigation  Collar drench	2L/ha  2 to 4L/ha	Apply in early Autumn or in Spring between 14 days prior to bud burst and full bloom. Late Summer or early Autumn application is recommended for early maturing cultivars. Divide rate per ha by the number of trees per ha to determine the rate per tree. Use high rate on medium and heavy soil types. <b>RETREATMENT:</b> Retreat trees with 2L/ha (Trickle) or with 2 or 3L/ha (Collar drench) in subsequent season. Rates of application for retreatment may be reduced if a lesser degree of growth control is desired. On light soils, retreat when normal growth resumes. Use the low rate for retreatment of low vigorous varieties in cool districts.
<b>Apricots Plums</b>	All States	Trickle irrigation  Collar drench	2L/ha  2 to 4L/ha	Apply in early Autumn or in Spring between 14 days prior to bud burst and full bloom. Bloom dates may be advanced. In plums, ensure that pollinator trees are treated at the same time to synchronise flowering. Divide rate per ha by the number of trees per ha to determine the rate per tree. Use high rate on medium and heavy soil types. <b>RETREATMENT:</b> Do not retreat until the season in which normal growth resumes.
<b>Cherries</b>	All States	Trickle irrigation  Collar Drench	1L/ha  1 to 2L/ha	Apply in early Autumn or in Spring between 14 days prior to bud burst and full bloom. Use high rate on medium and heavy soil types. Bloom dates may be advanced. Ensure that pollinator trees are treated at the same time to synchronise flowering. Divide rate per ha by the number of trees per ha to determine the rate per tree. <b>RETREATMENT:</b> Do not retreat until the season in which normal growth resumes.

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

## WITHHOLDING PERIOD

APPLES - DO NOT HARVEST FOR 21 DAYS AFTER APPLICATION.

## GENERAL INSTRUCTIONS

Payback is a systemic growth regulator that reduces internode lengths of new shoots and causes earlier formation of terminal buds. Use of Payback should begin only when trees are nearing their optimum size. It is most suited to intensively managed orchards where trees are near to filling their allocated space and growth control is a prime requirement. It is not suitable for old or unhealthy trees.

If a tree is under stress, Payback should NOT be used.

Stress can be caused by: moisture shortage due to lack of rain, or inadequate irrigation; high temperature; salinity; moisture excess due to water table, flooding or inappropriate irrigation; severe or successive frosts; poor fertility caused by imbalance of major or minor nutrients; diseases; insects; inappropriate herbicide use; mechanical damage.

## Tree Management

Treated trees may produce heavier crops and require a higher level of fertiliser application. Ensure that treated trees are not over-watered or subjected to drought stress. In some instances, treated trees may set excess fruit and careful attention should be given to thinning operations. When replanting a treated orchard, at least three years should elapse since the last application of Payback. In orchards which have been treated with collar drenches of Payback, fresh soil should be provided to allow tree establishment and growth. Remove treated soil. An alternative is to avoid original tree holes, and place young trees as far as practicable from the treated soil (minimum 1.5m).

Where young trees are required to be planted in an established orchard which has been treated with collar drenches of Payback, remove treated soil and replace with fresh soil.

## MIXING

**Foliar Sprays:** Half fill spray tank with clean water, add the required quantity of Payback and Shirwet 600 or Nufarm Surfactant 600 and agitate during the completion of filling. Frequent or constant agitation of the spray mixture is recommended. Wash out spray tank thoroughly after use.

**Collar drench:** Mix required volume of Payback with recommended volume of water. Periodic agitation may be necessary to prevent settling.

## APPLICATION

**Foliar sprays:** Apply in a sufficient volume of water to ensure penetration of the spray and good coverage of target tissue, new green stem tissue in the case of apples. Apply in a minimum volume of 1000L of water per ha. Shirwet 600 or Nufarm Surfactant 600 at 18 mL/100L of water must be added to the final spray volume. It is not normally necessary to reapply the spray if rain follows soon after application.

**Collar drench:** Up take of Payback is most efficient when it is applied to soil which receives adequate water and where there is a high density of feeder roots.

Apply required amount of Payback in 0.5 to 2 L of water per tree onto the soil/trunk crack line evenly around the base of the trunk. Where specific rates per tree are not given, the application rate per tree is determined by dividing the application rate per ha by the number of trees per ha. The volume of diluted drench required for even application without runoff will vary with the size of the tree and with soil type and moisture content. Use the higher volume on large trees.

Ensure that trees are adequately irrigated following treatment under dry conditions. Individual trees which are smaller than average should be left untreated or given a proportionally smaller dose of Payback.

**Trickle point:** Apply in 200-300 mL water per tree beneath each trickle or dripper point. Follow application with a short period of irrigation. Alternatively, the required amount of Payback can be applied through the trickle irrigation system. Equal distribution of water between trees is essential for this method to give a uniform effect.

## RETREATMENT

In some cases, control of growth may persist for more than one year. Retreat trees when normal growth resumes. Use the interval between the first treatment and the resumption of normal growth as a guide for retreatment in subsequent seasons.

## COMPATIBILITY

Payback is compatible with any one of the following products: Nimrod<sup>®</sup>, Omite<sup>®</sup>, Pirimor<sup>®</sup> or Dithane<sup>®</sup>. However, mixing Payback with more than any one of these products may give unpredictable results or may result in crop damage. If Carbaryl is used as a thinning spray on apples, allow 21 days before application of Payback.

## PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used container.

## STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well ventilated area. Do not store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

## SAFETY DIRECTIONS

Avoid contact with eyes and skin. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water.

## FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

## MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet.

### Conditions of sale

Crop Care Australasia Pty Ltd will not accept any responsibility whatsoever and howsoever arising and whether for consequential loss or otherwise in connection with the supply or use of these goods other than responsibility for the merchantable quality of the goods and such responsibilities mandatorily imposed by Statutes applicable to the sale or supply of these goods. To the extent allowed by such Statutes the liability of Crop Care Australasia Pty Ltd is limited to the replacement of the goods or (at the option of Crop Care Australasia Pty Ltd) the refund of the price paid and is conditional upon a claim being made in writing and where possible sufficient part of the goods to enable proper examination being returned to Crop Care Australasia Pty Ltd within thirty days of delivery.

In a Transport Emergency Dial 000 Police or Fire Brigade	SPECIALIST ADVICE IN EMERGENCY ONLY 1800 033 498 ALL HOURS – AUSTRALIA WIDE
--	---

Crop Care Australasia Pty Ltd is the licensed user of Payback which is a Registered Trademark. Shirwet is a Registered Trademark of Crop Care Australasia Pty Ltd. Nimrod, Pirimor, Omite and Dithane are Registered Trademarks.



Crop Care Australasia Pty Ltd  
ACN 061 362 347  
77 Tingira Street  
PINKENBA Qld 4008

Insert label code

APVMA Approval No. : 58627/0404